事 務 連 絡 令和4年9月14日

一般社団法人日本病院薬剤師会 御中

厚生労働省医薬・生活衛生局医薬品審査管理課

第十八改正日本薬局方(英文版)正誤表の送付について(その2)

標記について、別添写しのとおり各都道府県衛生主管部(局)薬務主管課宛に連絡しましたので、お知らせいたします。



事 務 連 絡 令和4年9月14日

各都道府県衛生主管部(局) 薬務主管課 御中

厚生労働省医薬・生活衛生局医薬品審査管理課

第十八改正日本薬局方(英文版)正誤表の送付について(その2)

第十八改正日本薬局方(令和3年厚生労働省告示第220号)の英文版につきまして、 一部に誤植等がありましたので別紙のとおり正誤表を送付いたします。

Official Monographs

Bicalutamide ビカルタミド

Page	Line	Correction	Error
550	left↑21-20	Determine each peak area by the automatic integration method: the peak areas of related substance M, having the relative retention time of about 0.26 to bicalutamide, related substance N, having the relative retention time of about 0.34, related substance K, having the relative retention time of about 1.03 and related substance L, having the relative retention time of about 1.13, obtained from the sample solution, are not larger than the peak area of bicalutamide from the standard solution,	Determine each peak area by the automatic integration method: the peak areas of related substance M, having the relative retention time of about 0.26 to bicalutamide, related substance N, having the relative retention time of about 0.34, related substance L, having the relative retention time of about 1.03 and related substance K, having the relative retention time of about 1.13, obtained from the sample solution, are not larger than the peak area of bicalutamide from the standard solution,

Candesartan Cilexetil and Amlodipine Besylate Tablets カンデサルタンシレキセチル・アムロジピンベシル酸塩錠

Page	Line	Correction	Error
615-618		Amlodipine Bes <u>i</u> late	Amlodipine Besylate

Imidapril Hydrochloride Tablets イミダプリル塩酸塩錠

The state of the s			
Page	Line	Correction	Error
1143	left ↑ 29-28	Add diluted methanol (2 in 5) to make 50 mL,	Add diluted ethanol (2 in 5) to make 50 mL,

Zopiclone ゾピクロン

Page	Line	Correction	Егтог
1935	right ↓ 33-36	determine each peak area by the automatic integration method: the peak areas of related substance A, having the relative retention time of about 0.1 to zopiclone, related substance B, having the relative retention time of about 0.2, related substance C, having the relative retention time of about 0.5, related substance D, having the relative retention time of about 0.9, obtained from the sample solution are not larger than 1/10 times the peak area of zopiclone from the standard solution, and the peaks mentioned above from the sample solution is not lager than 1/10 times the peak area of zopiclone from the standard solution.	determine each peak area by the automati integration method: the peak areas of related substance A, having the relative retention time of about 0.1 to zopiclone, related substance B having the relative retention time of about 0.2 related substance C, having the relative retention time of about 0.5, related substance D, having the relative retention time of about 0.9 and the peaks other than mentioned above obtained from the sample solution, are no larger than 1/10 times the peak area of zopiclone from the standard solution.